

From the bench to the bedside: ways to improve rituximab efficacy

Guillaume Cartron, Hervé Watier, Josée Golay, and Philippe Solal-Celigny

Rituximab (MabThera, Rituxan) is a chimeric IgG1 monoclonal antibody that specifically targets the CD20 surface antigen expressed on normal and neoplastic B-lymphoid cells. Rituximab is currently used in the treatment of both follicular and aggressive B-cell non-Hodgkin lymphomas. Despite its demonstrated clinical effectiveness, its in vivo mechanisms of action remain unknown and could differ by subtype of lymphoma. Rituximab has been shown to induce apoptosis, complement-mediated lysis, and antibody-dependent cellular cytotoxicity in vitro, and some evidence points toward an involvement of these mechanisms in vivo. Rituximab also has a delayed therapeutic effect as well as a potential “vaccinal” effect. Here, we review the current understanding of the mechanism of action of rituximab and discuss approaches that could increase its clinical activity. A better understanding of how rituximab acts in vivo should make it possible to develop new and more effective therapeutic strategies. (Blood. 2004;104:2635-2642)

fer by subtype of lymphoma. Rituximab has been shown to induce apoptosis, complement-mediated lysis, and antibody-dependent cellular cytotoxicity in vitro, and some evidence points toward an involvement of these mechanisms in vivo. Rituximab also has a delayed therapeutic effect as well as a potential “vaccinal” effect. Here, we review the current understanding of the mechanism of ac-

tion of rituximab and discuss approaches that could increase its clinical activity. A better understanding of how rituximab acts in vivo should make it possible to develop new and more effective therapeutic strategies. (Blood. 2004;104:2635-2642)

© 2004 by The American Society of Hematology

Introduction

Rituximab is a chimeric monoclonal antibody (mAb) produced by recombinant technology. It binds specifically to CD20, an antigen expressed by most human B lymphocytes. Rituximab was created by fusing the light- and heavy-chain variable domains of 2B8, a murine monoclonal anti-CD20 antibody and human κ light-chain and γ 1 heavy-chain constant regions.¹ Rituximab is currently indicated in both follicular and aggressive B-cell non-Hodgkin lymphomas (NHL). It is the first approved targeted treatment in the field of oncology and its impact on the treatment of NHL is evidenced by the short interval between its initial description¹ and its approval by both American and European authorities (1997 and 1998, respectively). Rituximab has some therapeutic activity in antibody-based autoimmune diseases,² and its indications are likely to be extended in the next few years. Rituximab has become the hallmark of anticancer mAbs, which offer a new approach to the treatment of B-cell lymphoproliferative disorders.

Although the clinical effectiveness of rituximab is no longer in question, its in vivo mechanisms of action have yet to be elucidated. In vitro data suggest that it induces apoptosis, complement-mediated lysis (complement-dependent cytotoxicity [CDC]), and antibody-dependent cellular cytotoxicity (ADCC) and some studies seem to confirm the involvement of these mechanisms in vivo. Other studies have suggested that rituximab may have a “vaccinal” effect. Most of these studies were performed with CD20⁺ cell lines in vitro and in murine models and in some cases with fresh neoplastic B cells in vitro. It is therefore difficult to evaluate the respective contribution of each of these effects that, moreover, could have different contributions according to lymphoma subtype. Several recent studies on CD20 structure and downstream signaling pathways have put forward key arguments that have helped to elucidate the in vivo mechanisms of action of

rituximab. A better understanding of rituximab-induced cell killing will allow the development of new, more effective antibodies derived from rituximab as well as innovative treatment modalities designed to optimize its mechanisms of action through selection of the most sensitive tumors, potentiation of cytotoxicity mechanisms or neutralization of resistance, design of optimal dosing schedules, and finding the best synergies. This review focuses on the current understanding of the mechanism of action of rituximab and suggests ways to improve its clinical efficacy.

The CD20 antigen

CD20 (or membrane-spanning 4-domain, group A, member 1 [MS4A1]) is a nonglycosylated protein of 33 to 35 kDa expressed on the surface of human B lymphocytes and also weakly on a small subset of T cells.³ The *MS4A1* gene is located on chromosome 11q12-13. It is switched on at the pre-B-cell stage of B-cell development, expressed throughout B-cell maturation, and lost during final maturation to plasma cells. Because CD20 antigen is not expressed by either plasma cells or B-lymphoid stem cells, rituximab does not affect significantly immunoglobulin serum concentrations, and after a conventional 4-dose course of rituximab therapy, B-lymphocyte counts recover in 9 to 12 months.^{4,5} The *MS4A1* promoter contains potential sites for several transcription factors such as Oct-1 and Oct-2, which bind to the BAT box,⁶ and PU.1 and Pip, which bind to the -160 site.⁷ Pip is a member of the interferon regulatory factor family and deletion of the BAT box and the -160 site results in an inactive promoter.^{6,8} Some evidence indicates that various cytokines including interferon α (IFN- α),

From the UPRES-EA Immuno-Pharmaco-Génétique des Anticorps thérapeutiques (IPGA), UPRES-EA, Université François Rabelais, Tours, France; Oncologie Médicale et Maladies du Sang; Laboratoire d'Immunologie, Centre Hospitalier Universitaire de Tours, France Laboratory of Cellular and Gene Therapy “G Lanzani,” Ospedali Riuniti Bergamo, Italia; Centre Jean Bernard, Le Mans, France; and Cancéropôle Grand Ouest, France.

Submitted March 24, 2004; accepted June 4, 2004. Prepublished online as *Blood* First Edition Paper, June 29, 2004; DOI 10.1182/blood-2004-03-1110.

Reprints: Guillaume Cartron, Oncologie Médicale et Maladies du Sang, CHRU Bretonneau, 2 bd Tonnelé, 37 044 Tours Cedex, France; e-mail: guillaume.cartron@med.univ-tours.fr.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 U.S.C. section 1734.

© 2004 by The American Society of Hematology

granulocyte-macrophage colony-stimulating factor (GM-CSF), interleukin-4 (IL-4), and tumor necrosis factor α (TNF- α) up-regulate CD20 expression on lymphoma cells in vitro and perhaps in vivo.⁹ However, the mechanism of this up-regulation is unknown. Although some of these cytokines may increase the therapeutic activity of rituximab,^{10,11} there is no direct evidence that this is related to increased CD20 expression.

The CD20 protein has 4 membrane-spanning domains with both the amino and carboxy termini of the protein located within the cytoplasm (Figure 1), and only a short extracellular segment of about 43 residues between the third and fourth transmembrane regions.¹² This configuration, deeply anchored in the membrane, protects from antigen shedding. Circulating "soluble forms" of CD20 have been detected in the serum of patients with chronic lymphocytic leukemia and high leukemic cell counts.¹³ These may originate from microparticles comprising CD20 inserted in their membrane bilayers. Human CD20 is 73% identical to murine CD20 with the transmembrane regions showing the greatest similarity of amino acid sequence.^{12,14} The extracellular domain differs in 16 of the approximate 43 amino acids,¹⁴ which may partly explain why rituximab does not bind to murine CD20. In the plasma membrane, the CD20 protein is probably not expressed in monomeric form, but organized into heterogeneous supramolecular complexes associating CD20 dimers or tetramers.¹⁵ CD20 has been reported to be closely associated with other proteins, in particular the transmembrane adapter protein p75/80, also named C-terminal src kinase-binding protein (Cbp),^{14,16,17} CD40, and major histocompatibility complex class II proteins (MHC II).¹⁸ Targeting CD20 with mAbs may therefore interfere with functions associated with CD20 partners such as CD40.¹⁸ Until recently, it was assumed that the extracellular loop bears only 2 groups of epitopes, one recognized by the majority of CD20 mAbs (such as B1, 2H7, rituximab), which induce inhibitory effects on B-cell proliferation, and another recognized only by the 1F5 mAb with unusual

activating properties.¹⁹⁻²³ However, Polyack et al¹⁴ have since demonstrated a high degree of epitope diversity among CD20 mAbs, which might explain the variety of biochemical and biologic effects of different anti-CD20 mAbs observed in vitro.²⁴ The extracellular alanine and proline at positions 170 and 172, respectively, are critical in determining the rituximab epitope, although neighboring residues as well as the overall state of the oligomeric complex seem to be important.

Antibody binding to CD20 induces rapid translocation of the molecule to lipid rafts,^{16,25} which are membrane microdomains enriched in cholesterol and sphingolipids (Figure 1). The association between CD20 and lipid rafts is dependent on cholesterol and on a short membrane-proximal cytoplasmic sequence (residues 219-225), which differs between murine and human CD20.²⁵ These lipid rafts are platforms for signal transduction, allowing the colocalization of receptors and signaling effectors. The likely association of CD20 with tyrosine kinases is supported by the observation that several of the biologic effects of anti-CD20 antibodies (homotypic aggregation, c-myc induction by 1F5 antibody) are dependent on tyrosine kinase.^{26,27} The presence of Cbp in association with CD20 in lipid rafts could concentrate a C-terminal Src kinase (Csk) such as Lyn, Fyn, or Lck,¹⁷ which phosphorylates Cbp, allowing autophosphorylation of Csk and thus inducing kinase activity.^{28,29} Recent data also suggest a role of acid sphingomyelinase and ceramide in CD20 signaling.³⁰ These recent studies have helped to elucidate CD20 antigen structure and localization in lipid rafts, providing insight into the diversity of anti-CD20 mAbs and their various biologic effects.

CD20 function and agonist effects of rituximab

No CD20 ligand has been described and CD20 does not display the usual structure of a receptor. CD20-deficient mice do not have any obvious defect of B-cell function.³¹ Therefore, the exact in vivo function of CD20 remains largely unknown. Recently, some data indicated that CD20 could play an important role in Ca²⁺ influx across plasma membranes, sustaining intracellular Ca²⁺ concentration and allowing activation of B cells.³² The usual, but artificial, technique to assess CD20-activated signaling events is based on the use of anti-CD20 agonist mAbs. From these studies, it appears that anti-CD20 mAbs regulate the cell cycle and also induce a wide variety of other biologic responses including an increase in the expression of MHC II or adhesion molecules (lymphocyte function-associated antigen [LFA-1, LFA-3]),²⁷ shedding of the B cell-expressed CD23,³³ down-regulation of the B-cell receptor,³⁴ and induction of apoptosis^{29,35-38} or, conversely, rescue from apoptosis.³⁹

The possibility that CD20 plays a crucial role in regulating cell cycle progression, at least in normal B cells, is supported by the fact that the mAb 1F5 induces cell cycle transition from G₀ to G₁, whereas B1-like mAbs inhibit B-cell progression from the G₁ to the S/G₂ plus M phases.¹⁹⁻²¹ Rituximab has been shown to induce apoptosis in numerous B-cell lines.^{29,36-38} However, induction of apoptosis in vivo frequently requires hypercross-linking, that is, cross-linking of antigen-bound anti-CD20 antibodies with secondary antibodies.^{29,36-38} Apoptosis is mediated by 2 main pathways leading to activation of cytosolic cysteine-proteases, named caspases (Figure 2). Intracellular stress signals induce apoptosis through the bcl-2 family of proteins and subsequent activation of caspase-9 (mitochondrial pathway), whereas the death-receptor pathway (Fas, TNF-related apoptosis-inducing ligand [TRAIL], and tumor




Figure 1. CD20 antigen structure and organization into membrane bilayers. (A) The CD20 protein has 4 cell membrane-spanning domains with both the amino and carboxy termini of the protein located within the cytoplasm. CD20 is closely associated with other proteins, in particular the C-terminal src kinase-binding protein (Cbp), CD40, and major histocompatibility complex class II proteins (MHC II). (B) Antibody binding to CD20 induces rapid translocation of the molecule to lipid rafts. Cbp can concentrate the C-terminal Src kinases (Csk) such as Lyn, Fyn, or Lck, which phosphorylate Cbp, leading to induction of kinase activity.




Figure 2. The 2 main apoptotic pathways. The “intrinsic” (mitochondrial) pathway is induced by intracellular stress signals leading to translocation of bcl-2 family members into the mitochondria. This induces alterations in mitochondrial permeability and consequent release of cytochrome C, which in association with apoptotic protease-activating factor 1 (APAF-1) activates caspase-9. The “extrinsic” (death-receptor) pathway is activated by the binding of ligands to tumor necrosis factor receptor (TNF-R) or FAS, inducing activation of caspase 8. Both caspase 8 and caspase-9 subsequently activate caspases-3, -6, and -7, which cleave substrates involved in the regulation of cell death.

necrosis factor [TNF] receptors) is induced by external signals and involves the activation of caspase-8. Caspases-8 and -9 then activate downstream caspases such as caspase-3, resulting in cell death. It has been reported that CD20-mediated Src activation leads to rapid phosphorylation of phospholipase C- γ 1 (PLC- γ 1) and PLC- γ 2,²⁶ which induces Ca²⁺ influx and caspase-3 activation.^{29,40} CD20-induced apoptosis is significantly reduced by Csk inhibitors,²⁹ caspase inhibitors, and calcium chelation,^{29,36,37} supporting the model in which CD20 ligation induces Csk activity, leading to PLC- γ phosphorylation, increased cytoplasmic Ca²⁺, and then caspase-3 activation.^{28,29} An important feature of apoptosis is its modulation by proapoptotic and antiapoptotic members of the bcl-2 gene family. The bcl-2 protein breaks the cell death machinery by blocking caspase activity, regulating Ca²⁺ flux, or by exerting an antioxidative activity.⁴¹⁻⁴³ Byrd et al⁴⁴ have demonstrated that a proportion of patients with B-cell chronic lymphocytic leukemia (B-CLL) treated with rituximab had in vivo activation of caspases-9 and -3, which correlated with depletion of B lymphocytes. This activation might be induced directly by rituximab but also by other mechanisms such as granzymes released by natural killer (NK) cells after binding of rituximab to Fc receptors.⁴⁵ Two recent studies have questioned the nature and role of caspases in apoptosis induction by anti-CD20 antibodies.^{46,47} These studies have shown that CD20-induced apoptosis is associated with membrane changes (phosphoserine translocation and mitochondrial permeability changes) but not with DNA fragmentation and chromatin condensation, unless CD20 hypercross-linking is present. Apoptosis is not blocked by caspase-specific concentrations of inhibitors and therefore appears caspase-independent. Moreover, bcl-2 overexpression blocks only mitochondrial permeability changes, but not apoptosis.^{46,47} These data are important in view of the role of bcl-2 in B-cell NHL, which overexpresses this protein, and considering the observation that rituximab associated with chemotherapy is able to overcome the adverse prognostic influence of bcl-2 overexpression by diffuse large B-cell lymphoma cells.⁴⁸

Recent clinical and in vitro data suggest that rituximab synergizes with chemotherapy by sensitizing malignant cells to the cytotoxic and apoptotic effects of drugs such as doxorubicin, cisplatin, dexamethasone, fludarabine, or retinoids.⁴⁹⁻⁵³ Rituximab

has also been reported to down-regulate IL-10 in some lymphoma cell lines. IL-10 normally promotes bcl-2 expression in B cells, making them more susceptible to apoptotic stimuli.^{54,55} The in vivo relevance of apoptosis induction by rituximab remains unclear. This antibody is a poor inducer of apoptosis in fresh normal, chronic lymphocytic leukemic, or B-lymphoma cells, in clear contrast to B-cell lines. Finally, 2B8-IgG4 mAb displays the same cross-linking as rituximab but has no effector functions and does not induce B-cell depletion in primates.⁵⁶ The in vivo reality of the apoptotic effects described for rituximab on cell lines still remains to be demonstrated in human lymphoma tumors.

CDC and CDCC

CDC is an important effector mechanism in the eradication of foreign agents and neoplastic cells. The complement system consists of classical, lectin, and alternative pathways that converge and ultimately generate the same set of effector molecules (Figure 3). The classical pathway requires immunoglobulins, whereas the lectin and alternative pathways are activated primarily by microbial components. The first step in activation of the classical pathway is the binding of the C1q component to Fc portions of IgG or IgM. This binding triggers a proteolytic cascade resulting in the generation of large amounts of C3b, the main effector molecule of the complement system. C3b molecules act as opsonins but also bind to the C3 convertase to form a C5 convertase, leading to the generation of the membrane attack complex (MAC) that kills target cells by disrupting the cell membrane. C3 is abundant in plasma and newly generated C3b is rapidly inactivated unless bound to the C5 convertase complex. C3b can also bind to complement receptors (CRs) expressed on effector cells such as granulocytes, macrophages, or NK cells and induce cell-mediated lysis or phagocytosis, depending on the effector cell complement-dependent cellular cytotoxicity (CDCC).

Several in vitro studies have demonstrated that rituximab induces CDC in B-lymphoma cell lines^{1,35,57} and fresh B-lymphoma cells,^{38,58-60} and this complement-mediated lysis correlates in part with the level of CD20 antigen expression.^{58,59,61} The sensitivity to CDC induced by rituximab could be different by subtype of lymphoma.^{59,61} Follicular and mantle cell lymphoma subtypes exhibited better responses to CDC than B-CLL.^{59,61} The ability of different anti-CD20 mAbs to induce complement-mediated lysis correlates with their capacity to translocate CD20 into lipid rafts,²⁴ an effect that could be dependent on the epitope




Figure 3. CDC classical pathway (IgG1, IgG3, IgM). The binding of the C1q component to Fc portions of IgG or IgM triggers a proteolytic cascade, resulting in the generation of large amounts of C3b. C3b molecules act as opsonins but also bind to the C3 convertase to form a C5 convertase, leading to the generation of the membrane attack complex (MAC), which kills the target.

recognized by the mAb. The increase in complement activation products observed during rituximab treatment⁶² and its correlation with the side effects observed after the first infusion⁶² confirm that complement is indeed activated *in vivo*, but evidence is still lacking in humans to establish a relationship between complement activation and therapeutic efficacy. The recent demonstration that rituximab is unable to cure C1q-deficient mice inoculated with syngenic lymphoma cells (EL4) transduced with human CD20 provides the first *in vivo* argument showing that complement activation is prerequisite to rituximab action.⁶³ Similar results have also been obtained in a complement-depleted xenograft model of lymphoma.⁶⁴ Finally, in agreement with the requirement for CDC and its dependence on CD20 expression levels is the observation that in cynomolgus monkeys, normal circulating B cells expressing low levels of CD20 antigen are more resistant than high expressers to rituximab treatment *in vivo*.⁶⁵

To prevent host tissue damage, a series of complement regulatory proteins (CtRPs) inhibits this system, among which glycosylphosphatidylinositol (GPI)-anchored proteins CD46, CD55, and CD59 are the most important (Figure 3). CD55 (decay-accelerating factor [DAF]) accelerates the decay of C3 and C5 convertase, and CD46 (membrane cofactor protein [MCP]) acts as a cofactor for the cleavage of C3b and C4b. CD59 prevents pore formation by the MAC. Convertase formation can also be prevented by cleaving C3b to its inactive form C3bi by factor I in conjunction with CD55 and CD46. The CtRPs are expressed by malignant B cells^{58,59} which might allow them to escape complement attack. Several reports have focused on CtRPs, particularly CD46, CD55, and CD59, in an attempt to relate their levels of expression with CDC. Although a correlation between CD20 expression level and CDC^{58,59} has been observed, the relationship between CD20/CtRP expression and rituximab-induced CDC has not been established.^{35,58-60,66} and the correlation with clinical response to rituximab still remains to be demonstrated.⁶⁰ Manches et al⁶¹ have recently shown that CDC *in vitro* correlates with the ratio of CD20 and CtRP intensities (CD46, CD55, CD59). The neutralization of CD59 and to some extent of CD55 and CD46 by blocking mAbs *in vitro* increases CDC.^{57,59,66} Therefore, concomitant targeting of CD20 and neutralization of CD59 by a bispecific mAb could be tested in rituximab-resistant B-cell lymphoid malignancies. Another way to act on CDC would be to improve the binding of C1q on the Fc portion. The C1q-binding site on human IgG1 has recently been mapped⁶⁷ and a mutant IgG1 with enhanced C1q binding has been created.⁶⁸ This mutant IgG1 has increased CDC activity but weakened ADCC *in vitro*.

Although rituximab is usually well tolerated, the first infusion may be complicated by severe first-dose side effects.⁴ These events are correlated with complement activation and with the number of circulating B cells.⁶² In some patients, rituximab infusion was associated with severe side effects described as “cytokine-release syndrome,”⁶⁹ which was fatal in a very limited number of cases.⁷⁰ However, the levels of cytokines assessed in these patients did not correlate with adverse events^{62,69} and complement activation could also be responsible for the severity of this side effect. Because complement activation is probably required for the therapeutic effect, the use of anticomplement medications should be avoided. Instead, a reduction of the infusion rate is usually effective in controlling “cytokine-release syndrome.”⁷¹ To summarize, *in vitro* and *in vivo* data suggest that CDC is an important mechanism underlying the action of rituximab and also some of its adverse reaction and future studies should be directed at determining its clinical significance as well as the

role of interactions between complement components and effector cells in rituximab activity.

Recruitment of Fc γ receptor-dependent functions

The CD20 protein is not shed from the cell surface and does not internalize on antibody binding and therefore appears to be an ideal target for the recruitment of effector cells expressing Fc γ receptors (Fc γ Rs). ADCC is an important effector mechanism in the eradication of intracellular pathogens and tumor cells. Binding of IgG to target cell antigens allows the recruitment of effector cells such as NK cells and macrophages, which express Fc receptors. These cells are then directed toward target cells, inducing either phagocytosis or release of their cytotoxic granules to promote cell killing. Immunoglobulins interact with effector cells through their Fc portion, which binds to specific receptors (Fc γ R) expressed by nearly all the cells of the immune system (Table 1). There are 3 classes of Fc γ R: Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). Fc γ RIa, Fc γ RIIa, Fc γ RIIc, and Fc γ RIIIa are activating receptors due to the presence of an immunoreceptor tyrosine-based activation motif (ITAM) either in the cytoplasmic domain (Fc γ RIIa, Fc γ RIIc) or in the accessory signaling γ chain (Fc γ RIa and Fc γ RIIIa). Cross-linking of activating Fc γ R by IgG-sensitized cells or IgG immune complexes induces tyrosine phosphorylation of ITAM by Src tyrosine kinases. Downstream signaling events eventually result in cellular responses such as phagocytosis, granule exocytosis, ADCC, or cytokine synthesis. Fc γ RIIb contains an immunoreceptor tyrosine-based inhibitory motif (ITIM) in its cytoplasmic domain and is the only inhibitory Fc receptor. Cocross-linking of Fc γ RIIb and BCR by an antigen already complexed to specific IgG results in tyrosine phosphorylation of ITIM, activation of phosphatases, and inhibition of BCR signals and B-cell activation. Most Fc γ Rs, including Fc γ RIIb, are expressed by macrophages. Therefore, immune complexes can engage activating and inhibitory Fc γ R on the same cell, providing an opportunity to fine-tune cell activation. NK cells only express the activating receptors, Fc γ RIIIa and Fc γ RIIc, whereas polymorphonuclear cells express Fc γ RIIa and Fc γ RIIIb, which synergize for phagocytosis.

In vitro studies have shown that rituximab induces ADCC in human lymphoma cell lines.^{1,35} In a xenograft model of lymphoma, the role of ADCC was clearly demonstrated using Fc γ - γ chain-deficient mice,⁷² which do not express activating Fc γ RI and Fc γ RIIIa, as well as in a model using a neutralizing antibody against murine Fc γ R.²² More recently, the role of both NK and polymorphonuclear neutrophils has been suggested in a disseminated lymphoma xenograft in severe combined immunodeficiency (SCID) mice, using antibody-dependent depletion of these cell

Table 1. Fc γ receptor expression according to cell type

	Fc γ RI	Fc γ RIIa	Fc γ RIIb	Fc γ RIIc	Fc γ RIIIa	Fc γ RIIIb
Monocytes/ macrophages	+	+	+	—	+	—
NK cells	—	—	—	+	+	—
Neutrophils	+/-	+	—	—	—	+
B lymphocytes	—	—	+	—	—	—
Dendritic cells	+	+	+	—	+	—
Mast cells	+/-	+	+	—	—	—
Platelets	—	+	—	—	—	—

types.⁷³ However, activating FcγRs are probably not the only receptors implicated in the antitumor response because FcγRIIb-deficient mice display a better tumoral reduction.⁷² These observations indicate that Fc receptor pathways are not restricted to an activating system such as expressed on NK cells, but also involve inhibitory pathways with FcγRIIb counterbalancing the activator receptors.

Human *FCGR3A* displays a nucleotide polymorphism at nucleotide 559 resulting in an amino acid substitution at position 158 with either a phenylalanine (F) or a valine (V). This substitution influences the affinity of IgG1 for the FcγRIIIa receptor with a higher affinity for FcγRIIIa-158VV NK cells.⁷⁴ We have demonstrated a correlation between the FcγRIIIa genotype and the clinical and molecular responses to rituximab in untreated follicular lymphoma; patients homozygous for FcγRIIIa-158VV have the best clinical and molecular responses.⁷⁵ These clinical results, recently confirmed by others,^{76,77} strongly support the hypothesis that ADCC is one of the in vivo mechanisms of rituximab action in follicular lymphoma. The lack of association in B-CLL⁷⁸ could suggest that the extent of ADCC in rituximab activity could be dependent of lymphoma proliferation subtype. Phagocytosis of human rituximab-opsonized lymphoma cells by macrophages has recently been demonstrated in vitro.⁶¹ This phagocytosis is mediated by FcγRIIa, indicating that other FcγR-expressing cells may be also involved in the action of rituximab.⁶¹

The importance of the interaction between the Fc portion and FcγR highlights a potential way to increase rituximab activity. By in vitro molecular engineering of the Fc region, it is possible to improve the affinity of mAbs for FcγRIIIa and thereby to enhance ADCC.⁷⁹ Glycosylation of IgG is necessary for effector functions and particularly for the interaction with FcγR.⁸⁰ Modifications of glycosylation could influence the ability of mAbs to activate ADCC. For instance, addition of bisecting *N*-acetylglucosamine or a lack of fucose on IgG1 *N*-linked oligosaccharide chain can dramatically increase ADCC.⁸¹⁻⁸³

As shown in the pivotal study,^{4,84} the pharmacokinetic profile of rituximab differs between responders and nonresponders. Moreover, normal B-cell depletion appears to correlate with circulating levels of rituximab.⁸⁵ It has been suggested that the *FCGR3A*

genotype affects rituximab pharmacokinetics, but this requires confirmation in a larger number of patients.

Cytokines such as IL-2, IL-12, IFN-α, GM-CSF, and granulocyte colony-stimulating factor (G-CSF) are known to increase ADCC/phagocytosis by stimulation or expansion of NK cells and macrophages.⁸⁶⁻⁸⁹ For example, IL-2 increases ADCC by NK cell expansion and activation⁸⁸ and a correlation between NK cell counts and clinical response was found in a clinical trial combining IL-2 and rituximab in patients with NHL.⁹⁰ IL-12 enhances the lytic activity of NK cells and induces IFN-γ.⁸⁹ This cytokine was recently used in combination with rituximab in patients with recurrent NHL with a promising objective response rate.⁹¹ IFN-α has also been used in combination with rituximab in patients with relapsed NHL. The response rates were similar to those observed in trials where rituximab was given alone, although IFN-α increased time to treatment failure in responding patients.^{10,92} GM-CSF induces monocyte differentiation and can activate polynuclear neutrophils.⁸⁶ The combination of rituximab with GM-CSF has been used in patients with relapsed/refractory NHL and preliminary results suggested a complete remission rate higher than that in patients treated with rituximab alone.¹¹ In patients who received high-dose therapy and an autologous stem cell transplant, the period after grafting, which is associated with rapid NK cell count recovery,⁹³ could be an optimal window for rituximab administration.⁹⁴ In contrast, steroids have a dual impact on lymphoma cell lines. They increase apoptosis of lymphoma lines but also lead to decreased ADCC.⁵² Thus, the available evidence strongly favors a role for ADCC in the clinical activity of rituximab, paving the way to future studies to better define strategies for maximizing ADCC through the use of anti-CD20 mAbs.

Other mechanisms of action and “vaccinal” effect of rituximab

Several studies have indicated that maximal clinical and molecular responses to rituximab therapy may take several months, suggesting that short-term cytolytic mechanisms such as apoptosis, CDC, and ADCC are not the only ones involved (Figure 4). Rituximab-




Figure 4. Main mechanisms of action of rituximab and ways to increase its clinical efficacy. FAMP indicates fludarabine monophosphate; CR, complement receptor.

promoted lysis of lymphoma cells through any of these latter mechanisms may promote uptake and cross-presentation of lymphoma cell–derived peptides by dendritic cells, inducing their maturation and allowing the generation of specific cytotoxic T lymphocytes.⁹⁵ It is difficult to demonstrate such an immune reaction in humans, but murine experiments clearly indicate that passive immunotherapy with mAbs can induce a specific antitumor immunity^{96,97} (not directed against the target antigen but against several unknown tumor-derived antigens). Induction of this specific immunity requires myeloid-derived dendritic cells,^{97,98} suggesting that adjuvant therapy promoting this pathway (GM-CSF, Flt-3 ligand, and IFN- α) might be of benefit. In this context, it is noteworthy that the combination of rituximab with GM-CSF yielded promising results in a phase 2 trial.¹¹ Clinical studies have shown that retreatment with rituximab in patients who had a relapse after an initial response increased the time to progression⁹⁹ and that maintenance therapy given to patients in response to rituximab increased the response rates.¹⁰⁰ Together, these results strongly suggest that rituximab may have a “vaccinal” effect and may therefore be useful as maintenance therapy. Randomized clinical trials are needed to confirm the clinical impact of this approach.

Conclusion

A large body of evidence shows that rituximab induces cell death by different pathways, in particular ADCC, CDC, and apoptosis.

References

- Reff ME, Carner K, Chambers KS, et al. Depletion of B cells in vivo by a chimeric mouse monoclonal antibody to CD20. *Blood*. 1994;83:435-445.
- Edwards JC, Cambridge G. Sustained improvement in rheumatoid arthritis following a protocol designed to deplete B lymphocytes. *Rheumatology (Oxford)*. 2001;40:205-211.
- Hultin LE, Hausner MA, Hultin PM, Giorgi JV. CD20 (pan-B cell) antigen is expressed at a low level on a subpopulation of human T lymphocytes. *Cytometry*. 1993;14:196-204.
- McLaughlin P, Grillo-Lopez AJ, Link BK, et al. Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a four-dose treatment program. *J Clin Oncol*. 1998;16:2825-2833.
- Maloney DG, Grillo-Lopez AJ, Bodkin DJ, et al. IDEC-C2B8: results of a phase I multiple-dose trial in patients with relapsed non-Hodgkin's lymphoma. *J Clin Oncol*. 1997;15:3266-3274.
- Thevenin C, Lucas BP, Kozlow EJ, Kehrl JH. Cell type- and stage-specific expression of the CD20/B1 antigen correlates with the activity of a diverged octamer DNA motif present in its promoter. *J Biol Chem*. 1993;268:5949-5956.
- Hromas R, Orazi A, Neiman RS et al. Hematopoietic lineage- and stage-restricted expression of the ETS oncogene family member PU.1. *Blood*. 1993;82:2998-3004.
- Himmelfmann A, Riva A, Wilson GL, Lucas BP, Thevenin C, Kehrl JH. PU.1/Pip and basic helix loop helix zipper transcription factors interact with binding sites in the CD20 promoter to help confer lineage- and stage-specific expression of CD20 in B lymphocytes. *Blood*. 1997;90:3984-3995.
- Venugopal P, Sivaraman S, Huang XK, Nayini J, Gregory SA, Preisler HD. Effects of cytokines on CD20 antigen expression on tumor cells from patients with chronic lymphocytic leukemia. *Leuk Res*. 2000;24:411-415.
- Davis TA, Maloney DG, Grillo-Lopez AJ, White CA, et al. Combination immunotherapy of relapsed or refractory low-grade or follicular non-Hodgkin's lymphoma with rituximab and interferon- α -2a. *Clin Cancer Res*. 2000;6:2644-2652.
- Rossi JF, Yang LZ, Quittet P, et al. Rituximab and GM-CSF, an effective therapy for relapsed/refractory patients with low-grade B-cell lymphoma: correlation between response and dendritic cell subpopulation mobilized [abstract]. *Blood*. 2001;98:607a.
- Einfeld DA, Brown JP, Valentine MA, Clark EA, Ledbetter JA. Molecular cloning of the human B cell CD20 receptor predicts a hydrophobic protein with multiple transmembrane domains. *EMBO J*. 1988;7:711-717.
- Manshour T, Do KA, Wang X, et al. Circulating CD20 is detectable in the plasma of patients with chronic lymphocytic leukemia and is of prognostic significance. *Blood*. 2003;101:2507-2513.
- Polyak MJ, Deans JP. Alanine-170 and proline-172 are critical determinants for extracellular CD20 epitopes; heterogeneity in the fine specificity of CD20 monoclonal antibodies is defined by additional requirements imposed by both amino acid sequence and quaternary structure. *Blood*. 2002;99:3256-3262.
- Bubien JK, Zhou LJ, Bell PD, Frizzell RA, Tedder TF. Transfection of the CD20 cell surface molecule into ectopic cell types generates a Ca²⁺ conductance found constitutively in B lymphocytes. *J Cell Biol*. 1993;121:1121-1132.
- Deans JP, Li H, Polyak MJ. CD20-mediated apoptosis: signalling through lipid rafts. *Immunology*. 2002;107:176-182.
- Deans JP, Kalt L, Ledbetter JA, Schieven GL, Bolen JB, Johnson P. Association of 75/80-kDa phosphoproteins and the tyrosine kinases Lyn, Fyn, and Lck with the B cell molecule CD20. Evidence against involvement of the cytoplasmic regions of CD20. *J Biol Chem*. 1995;270:22632-22638.
- Leveille C, Al-Daccak R, Mourad W. CD20 is physically and functionally coupled to MHC class II and CD40 on human B cell lines. *Eur J Immunol*. 1999;29:65-74.
- Clark EA, Shu G, Ledbetter JA. Role of the Bp35 cell surface polypeptide in human B-cell activation. *Proc Natl Acad Sci U S A*. 1985;82:1766-1770.
- Golay JT, Clark EA, Beverley PC. The CD20 (Bp35) antigen is involved in activation of B cells from the G0 to the G1 phase of the cell cycle. *J Immunol*. 1985;135:3795-3801.
- Smeland EB, Beiske K, Ohlsson R, et al. Activation of human B cells: alternate options for initial triggering and effects of nonmitogenic concentrations of anti-IgM antibodies on resting and activated cells. *J Immunol*. 1987;138:3179-3184.
- Funakoshi S, Longo DL, Murphy WJ. Differential in vitro and in vivo antitumor effects mediated by anti-CD40 and anti-CD20 monoclonal antibodies against human B-cell lymphomas. *J Immunother Emphasis Tumor Immunol*. 1996;19:93-101.
- Press OW, Appelbaum F, Ledbetter JA, et al. Monoclonal antibody 1F5 (anti-CD20) serotherapy of human B cell lymphomas. *Blood*. 1987;69:584-591.
- Cragg MS, Morgan SM, Chan HTC, et al. Complement-mediated lysis by anti-CD20 mAb correlates with segregation into lipid rafts. *Blood*. 2003;101:1045-1052.
- Polyak MJ, Taylor SH, Deans JP. Identification of a cytoplasmic region of CD20 required for its redistribution to a detergent-insoluble membrane compartment. *J Immunol*. 1998;161:3242-3248.
- Deans JP, Schieven GL, Shu GL, et al. Association of tyrosine and serine kinases with the B cell surface antigen CD20. Induction via CD20 of tyrosine phosphorylation and activation of phospholipase C- γ 1 and PLC phospholipase C- γ 2. *J Immunol*. 1993;151:4494-4504.
- White MW, McConnell F, Shu GL, Morris DR, Clark EA. Activation of dense human tonsillar B cells. Induction of c-myc gene expression via two

Acknowledgments

We wish to thank the CANCEC Association for its action in improving patient hospitalization conditions and their financial support in cancer research.

- distinct signal transduction pathways. *J Immunol.* 1991;146:846-853.
28. Semac I, Palomba C, Kulangara K, et al. Anti-CD20 therapeutic antibody rituximab modifies the functional organization of rafts/microdomains of B lymphoma cells. *Cancer Res.* 2003;63:534-540.
 29. Hofmeister JK, Cooney D, Coggeshall KM. Clustered CD20 induced apoptosis: src-family kinase, the proximal regulator of tyrosine phosphorylation, calcium influx, and caspase 3-dependent apoptosis. *Blood Cells Mol Dis.* 2000;26:133-143.
 30. Bezombes C, Graziade S, Garret C, et al. Rituximab antiproliferative effect in B lymphoma cells is associated with acid-sphingomyelinase activation in raft microdomains. *Blood.* 2004;104:1166-1173.
 31. O'Keefe TL, Williams GT, Davies SL, Neuberger MS. Mice carrying a CD20 gene disruption. *Immunogenetics.* 1998;48:125-132.
 32. Li H, Ayert LM, Lytton J, Deans JP. Store-operated cation entry mediated by CD20 in membrane rafts. *J Biol Chem.* 2003;278:42427-42434.
 33. Bourget I, Di Berardino W, Breittmayer JP, et al. CD20 monoclonal antibodies stimulate extracellular cleavage of the low affinity receptor for IgE (Fc epsilon RII/CD23) in Epstein-Barr-transformed B cells. *J Biol Chem.* 1994;269:6927-6930.
 34. Bourget I, Breittmayer JP, Grenier-Brossette N, Cousin JL. CD20 monoclonal antibodies down-regulate IgM at the surface of B cells. *Eur J Immunol.* 1993;23:768-771.
 35. Fliedger D, Renoth S, Beier I, Sauerbruch T, Schmidt-Wolf I. Mechanism of cytotoxicity induced by chimeric mouse human monoclonal antibody IDEC-C2B8 in CD20-expressing lymphoma cell lines. *Cell Immunol.* 2000;204:55-63.
 36. Shan D, Ledbetter JA, Press OW. Signaling events involved in anti-CD20-induced apoptosis of malignant human B cells. *Cancer Immunol Immunother.* 2000;48:673-683.
 37. Mathas S, Rickers A, Bommert K, Dorken B, Mapara MY. Anti-CD20- and B-cell receptor-mediated apoptosis: evidence for shared intracellular signaling pathways. *Cancer Res.* 2000;60:7170-7176.
 38. Golay J, Zaffaroni L, Vaccari T, et al. Biologic response of B lymphoma cells to anti-CD20 monoclonal antibody rituximab in vitro: CD55 and CD59 regulate complement-mediated cell lysis. *Blood.* 2000;95:3900-3908.
 39. Valentine MA, Licciardi KA. Rescue from anti-IgM-induced programmed cell death by the B cell surface proteins CD20 and CD40. *Eur J Immunol.* 1992;22:3141-3148.
 40. Shan D, Ledbetter JA, Press OW. Apoptosis of malignant human B cells by ligation of CD20 with monoclonal antibodies. *Blood.* 1998;91:1644-1652.
 41. Genestier L, Bonnefoy-Berard N, Rouault JP, Flacher M, Revillard JP. Tumor necrosis factor-alpha up-regulates Bcl-2 expression and decreases calcium-dependent apoptosis in human B cell lines. *Int Immunol.* 1995;7:533-540.
 42. Hockenbery DM, Oltvai ZN, Yin XM, Millman CL, Korsmeyer SJ. Bcl-2 functions in an antioxidant pathway to prevent apoptosis. *Cell.* 1993;75:241-251.
 43. Baffy G, Miyashita T, Williamson JR, Reed JC. Apoptosis induced by withdrawal of interleukin-3 (IL-3) from an IL-3-dependent hematopoietic cell line is associated with repartitioning of intracellular calcium and is blocked by enforced Bcl-2 oncoprotein production. *J Biol Chem.* 1993;268:6511-6519.
 44. Byrd JC, Kitada S, Flinn IW, et al. The mechanism of tumor cell clearance by rituximab in vivo in patients with B-cell chronic lymphocytic leukemia: evidence of caspase activation and apoptosis induction. *Blood.* 2002;99:1038-1043.
 45. Lieberman J. The ABCS of granule-mediated cytotoxicity: new weapons in the arsenal. *Nat Rev Immunol.* 2003;3:361-370.
 46. Chan HT, Hughes D, French RR, et al. CD20-induced lymphoma cell death is independent of both caspases and its redistribution into triton X-100 insoluble membrane rafts. *Cancer Res.* 2003;63:5480-5489.
 47. van der Kolk LE, Evers LM, Omene C, et al. CD20-induced B cell death can bypass mitochondria and caspase activation. *Leukemia.* 2002;16:1735-1744.
 48. Mounier N, Briere J, Gisselbrecht C, et al. Rituximab plus CHOP (R-CHOP) overcomes bcl-2-associated resistance to chemotherapy in elderly patients with diffuse large B-cell lymphoma (DLBCL). *Blood.* 2003;101:4279-4284.
 49. Shan D, Gopal AK, Press OW. Synergistic effects of the fenretinide (4-HPR) and anti-CD20 monoclonal antibodies on apoptosis induction of malignant human B cells. *Clin Cancer Res.* 2001;7:2490-2495.
 50. Ghetie MA, Bright H, Vitetta ES. Homodimers but not monomers of Rituxan (chimeric anti-CD20) induce apoptosis in human B-lymphoma cells and synergize with a chemotherapeutic agent and an immunotoxin. *Blood.* 2001;97:1392-1398.
 51. Di Gaetano N, Xiao Y, Erba E, et al. Synergism between fludarabine and rituximab revealed in a follicular lymphoma cell line resistant to the cytotoxic activity of either drug alone. *Br J Haematol.* 2001;114:800-809.
 52. Rose AL, Smith BE, Maloney DG. Glucocorticoids and rituximab in vitro: synergistic direct antiproliferative and apoptotic effects. *Blood.* 2002;100:1765-1773.
 53. Alas S, Ng CP, Bonavida B. Rituximab modifies the cisplatin-mitochondrial signaling pathway, resulting in apoptosis in cisplatin-resistant non-Hodgkin's lymphoma. *Clin Cancer Res.* 2002;8:836-845.
 54. Alas S, Emmanouilides C, Bonavida B. Inhibition of interleukin 10 by rituximab results in down-regulation of bcl-2 and sensitization of B-cell non-Hodgkin's lymphoma to apoptosis. *Clin Cancer Res.* 2001;7:709-723.
 55. Alas S, Bonavida B. Rituximab inactivates signal transducer and activation of transcription 3 (STAT3) activity in B-cell non-Hodgkin's lymphoma through inhibition of the interleukin 10 autocrine/paracrine loop and results in down-regulation of Bcl-2 and sensitization to cytotoxic drugs. *Cancer Res.* 2001;61:5137-5144.
 56. Anderson DR, Grillo-Lopez AJ, Varns C, Chambers KS, Hanna N. Targeted anti-cancer therapy using rituximab, a chimaeric anti-CD20 antibody (IDEC-C2B8) in the treatment of non-Hodgkin's B-cell lymphoma. *Biochem Soc Trans.* 1997;25:705-708.
 57. Harjunpaa A, Junnikkala S, Meri S. Rituximab (anti-CD20) therapy of B-cell lymphomas: direct complement killing is superior to cellular effector mechanisms. *Scand J Immunol.* 2000;51:634-641.
 58. Golay J, Lazzari M, Facchinetti V, et al. CD20 levels determine the in vitro susceptibility to rituximab and complement of B-cell chronic lymphocytic leukemia: further regulation by CD55 and CD59. *Blood.* 2001;98:3383-3389.
 59. Bellosillo B, Villamor N, Lopez-Guillermo A, et al. Complement-mediated cell death induced by rituximab in B-cell lymphoproliferative disorders is mediated in vitro by a caspase-independent mechanism involving the generation of reactive oxygen species. *Blood.* 2001;98:2771-2777.
 60. Weng WK, Levy R. Expression of complement inhibitors CD46, CD55, and CD59 on tumor cells does not predict clinical outcome after rituximab treatment in follicular non-Hodgkin lymphoma. *Blood.* 2001;98:1352-1357.
 61. Manches O, Lui G, Chaperot L, et al. In vitro mechanisms of action of rituximab on primary non-Hodgkin lymphomas. *Blood.* 2003;101:949-954.
 62. van der Kolk LE, Grillo-Lopez AJ, Baars JW, Hack CE, van Oers MH. Complement activation plays a key role in the side-effects of rituximab treatment. *Br J Haematol.* 2001;115:807-811.
 63. Di Gaetano N, Cittera E, Nota R, et al. Complement activation determines the therapeutic activity of rituximab in vivo. *J Immunol.* 2003;171:1581-1587.
 64. Cragg MS, Glennie MJ. Antibody specificity controls in vivo effector mechanisms of anti-CD20 reagents. *Blood.* 2004;103:2738-2743.
 65. Vugmeyster Y, Howell K, Bakshi A, Flores C, Canova-Davis E. Effect of anti-CD20 monoclonal antibody, Rituxan, on cynomolgus monkey and human B cells in a whole blood matrix. *Cytometry.* 2003;52A:101-109.
 66. Treon SP, Mitsiades C, Mitsiades N, et al. Tumor cell expression of CD59 is associated with resistance to CD20 serotherapy in patients with B-cell malignancies. *J Immunother.* 2001;24:263-271.
 67. Idusogie EE, Presta LG, Gazzano-Santoro H, et al. Mapping of the C1q binding site on rituxan, a chimeric antibody with a human IgG1 Fc. *J Immunol.* 2000;164:4178-4184.
 68. Idusogie EE, Wong PY, Presta LG, et al. Engineered antibodies with increased activity to recruit complement. *J Immunol.* 2001;166:2571-2575.
 69. Winkler U, Jensen M, Manzke O, Schulz H, Diehl V, Engert A. Cytokine-release syndrome in patients with B-cell chronic lymphocytic leukemia and high lymphocyte counts after treatment with an anti-CD20 monoclonal antibody (rituximab, IDEC-C2B8). *Blood.* 1999;94:2217-2224.
 70. Lim LC, Koh LP, Tan P. Fatal cytokine release syndrome with chimeric anti-CD20 monoclonal antibody rituximab in a 71-year-old patient with chronic lymphocytic leukemia. *J Clin Oncol.* 1999;17:1962-1963.
 71. Buysmann S, Hack CE, van Diepen FN, Surachno J, ten Berge IJ. Administration of OKT3 as a two-hour infusion attenuates first-dose side effects. *Transplantation.* 1997;64:1620-1623.
 72. Clynes RA, Towers TL, Presta LG, Ravetch JV. Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. *Nat Med.* 2000;6:443-446.
 73. Hernandez-Illizaliturri FJ, Jupudy V, Ostberg J, et al. Neutrophils contribute to the biological antitumor activity of rituximab in a non-Hodgkin's lymphoma severe combined immunodeficiency mouse model. *Clin Cancer Res.* 2003;9:5866-5873.
 74. Koene HR, Kleijer M, Algra J, Roos D, von dem Borne AE, de Haas M. Fc gammaRIIIa-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gammaRIIIa, independently of the Fc gammaRIIIa-48L/R/H phenotype. *Blood.* 1997;90:1109-1114.
 75. Cartron G, Dacheux L, Salles G, et al. Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor FcγRIIIa gene. *Blood.* 2002;98:754-758.
 76. Weng WK, Levy R. Two immunoglobulin G fragment C receptor polymorphisms independently predict response to rituximab in patients with follicular lymphoma. *J Clin Oncol.* 2003;21:3940-3947.
 77. Weng WK, Levy R. Analysis of IgG Fc receptor FcγRIIIa polymorphism in relapsed follicular non-Hodgkin's lymphoma patients treated with rituximab [abstract]. *Blood.* 2002;100:1368a.
 78. Farag SS, Flinn IW, Lehman TA, Young D, Byrd JC. FcγRIIIa and FcγRIIIa polymorphisms do not predict response to rituximab in B-cell chronic lymphocytic leukemia. *Blood.* 2004;103:1472-1474.
 79. Shields RL, Namenuk AK, Hong K, et al. High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma

- RIII, and FcRn and design of IgG1 variants with improved binding to the Fc gamma R. *J Biol Chem*. 2001;276:6591-6604.
80. Wright A, Morrison SL. Effect of glycosylation on antibody function: implications for genetic engineering. *Trends Biotechnol*. 1997;15:26-32.
81. Umama P, Jean-Mairet J, Moudry R, Amstutz H, Bailey JE. Engineered glycoforms of an antineuroblastoma IgG1 with optimized antibody-dependent cellular cytotoxic activity. *Nat Biotechnol*. 1999;17:176-180.
82. Shinkawa T, Nakamura K, Yamane N, et al. The absence of fucose but not the presence of galactose or bisecting N-acetylglucosamine of human IgG1 complex-type oligosaccharides shows the critical role enhancing antibody-dependent cellular toxicity. *J Biol Chem*. 2003;278:3466-3473.
83. Davies J, Jiang L, Pan LZ, LaBarre MJ, Anderson D, Reff M. Expression of GnTIII in a recombinant anti-CD20 CHO production cell line: Expression of antibodies with altered glycoforms leads to an increase in ADCC through higher affinity for Fc-gammaRIII. *Biotechnol Bioeng*. 2001;74:288-294.
84. Berinstein NL, Grillo-Lopez AJ, White CA, et al. Association of serum rituximab (IDEC-C2B8) concentration and anti-tumor response in the treatment of recurrent low-grade or follicular non-Hodgkin's lymphoma. *Ann Oncol*. 1998;9:995-1001.
85. Anolik JH, Campbell D, Felgar RE, et al. The relationship of Fc-gammaRIIIa genotype to degree of B cell depletion by rituximab in the treatment of systemic lupus erythematosus. *Arthritis Rheum*. 2003;48:455-459.
86. Wing EJ, Magee DM, Whiteside TL, Kaplan SS, Shadduck RK. Recombinant human granulocyte/macrophage colony-stimulating factor enhances monocyte cytotoxicity and secretion of tumor necrosis factor alpha and interferon in cancer patients. *Blood*. 1989;73:643-646.
87. Herberman RB, Ortaldo JR, Mantovani A, Hobbs DS, Kung HF, Pestka S. Effect of human recombinant interferon on cytotoxic activity of natural killer (NK) cells and monocytes. *Cell Immunol*. 1982;67:160-167.
88. Phillips JH, Lanier LL. Dissection of the lymphokine-activated killer phenomenon. Relative contribution of peripheral blood natural killer cells and T lymphocytes to cytotoxicity. *J Exp Med*. 1986;164:814-825.
89. Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol*. 2003;3:133-146.
90. Wilson S, Hurst D, Yuen A, et al. IL-2 mediated NK cell expansion correlates with clinical response to rituximab: results of two phase I trials of the combination of IL-2 and rituximab [abstract]. *Blood*. 2001;98:602a.
91. Ansell SM, Witzig TE, Kurtin PJ, et al. Phase 1 study of interleukin-12 in combination with rituximab in patients with B-cell non-Hodgkin lymphoma. *Blood*. 2002;99:67-74.
92. Sacchi S, Federico M, Vitolo U, et al. Clinical activity and safety of combination immunotherapy with IFN-alpha 2a and rituximab in patients with relapsed low grade non-Hodgkin's lymphoma. *Haematologica*. 2001;86:951-958.
93. Guillaume T, Rubinstein DB, Symann M. Immune reconstitution and immunotherapy after autologous hematopoietic stem cell transplantation. *Blood*. 1998;92:1471-1490.
94. Shimoni A, Hardan I, Avigdor A, Yeshurun M, Raanani P. Rituximab reduces relapse risk after allogeneic and autologous stem cell transplantation in patients with high-risk aggressive non-Hodgkin's lymphoma. *Br J Haematol*. 2003;122:457-464.
95. Selenko N, Majdic O, Jager U, Sillaber C, Stockl J, Knapp W. Cross-priming of cytotoxic T cells promoted by apoptosis-inducing tumor cell reactive antibodies? *J Clin Immunol*. 2002;22:124-130.
96. Rafiq K, Bergtold A, Clynes R. Immune-complex-mediated antigen presentation induces tumor immunity. *J Clin Invest*. 2002;110:71-79.
97. Kalergis AM, Ravetch JV. Inducing tumor immunity through the selective engagement of activating Fc-gamma receptors on dendritic cells. *J Exp Med*. 2002;195:1653-1659.
98. Dyall R, Vasovic LV, Clynes R, Nikolic-Zugic J. Cellular requirements for the monoclonal antibody-mediated eradication of an established solid tumor. *Eur J Immunol*. 1999;29:30-37.
99. Davis TA, Grillo-Lopez AJ, White CA, et al. Rituximab anti-CD20 monoclonal antibody therapy in non-Hodgkin's lymphoma: safety and efficacy of re-treatment. *J Clin Oncol*. 2000;18:3135-3143.
100. Hainsworth JD, Litchy S, Burris HA, Scullin DC, Corso SW, Yardley DA, Moriss Greco FA. Rituximab as first-line maintenance therapy for patients with indolent non-lymphoma. *J Clin Oncol*. 2002;20:4261-4267.